

Entry of this amendment is requested.

Respectfully submitted,

OBLON, SPIVAK, McCLELLAND,  
MAIER & NEUSTADT, P.C.

Norman F. Oblon  
Attorney of Record  
Registration No. 24,618

Samuel H. Blech  
Registration No. 32,082



**22850**


(703) 413-3000  
Fax No.: (703) 413-2220  
NFO:SHB/cja  
I:\atty\SHB\106440001-am2.wpd

**Marked-Up Copy**  
Serial No: 09/445,439  
Amendment Filed Herewith

IN THE CLAIMS



--101. (Twice Amended) A method for preparing a drug targeting system for administering one or more physiologically effective substances to a mammal, said method comprising:

a) preparing nanoparticles made of a polymeric material, said nanoparticles being free of a surfactant surface coating and comprising said polymeric material, one or more physiologically effective substances to be delivered to said mammal, and one or more stabilizers for said nanoparticles allowing targeting of said one or more physiologically effective substances to a specific site within or on a mammalian body, said one or more stabilizers being selected from the group consisting of polysorbate 85, polysorbate 81, dextran 12,000, carboxylic acid esters of glycerol, sorbitan monostearate, sorbitan monooleate, polyoxamer 188, polyoxamines, alkoxyated ethers, alkoxyated esters, alkoxyated mono-, di and triglycerides, alkoxyated phenols and diphenols, Genapol® compounds, Bauki compounds<sup>R</sup>, sodium stearate, metal salts of carboxylic acids, metal salts of alcohol sulfates and metal salts of sulfosuccinates and mixtures of two or more of said substances, wherein said Genapol<sup>R</sup> compounds are of the formula



wherein y is in the range of 4 to 18 and x is in the range of 1 to 18,

and said Bauki<sup>R</sup> compounds are of the formulas (I) or (I')



in which  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_6$  are identical or different and represent hydrogen and a methyl or ethyl group,

$Q$  represents a valency, oxygen or an ester or amide bridge and  $Q'$  denotes hydrogen if  $Q$  represents a valency or oxygen, and is a hydroxyl or amino group if  $Q$  represents an ester or amide bridge,

$x$  is an integer from 3 to 50, if  $Q$  is a valency or oxygen, and an integer from 3 to 1000, if  $Q$  is an ester or amide function,  $G_1$  and  $G_2$  are a valency, oxygen or an ester or amide group, it being possible for the two groups to be identical or different,  $n$  is an integer from 4 to 44,  $y$  is an integer from 2 to 50, and  $R_3$  is hydrogen or a lower alkyl having 1-6 C atoms, and; by polymerizing one or more monomeric or oligomeric precursors of said polymeric material or both, in the presence of said one or more physiologically effective substances and in the presence of said stabilizers; and optionally

b) providing said nanoparticles in a medium allowing the transport of said nanoparticles to a target within or on said mammal after administration.--